

## Review Article

## The sequel to COVID-19: the antithesis to life

Sudhir Bhandari<sup>1</sup>, Ajit Singh Shaktawat<sup>1</sup>, Bhoopendra Patel<sup>2\*</sup>, Amitabh Dube<sup>3</sup>, Shivankan Kakkar<sup>4</sup>, Amit Tak<sup>3</sup>, Jitendra Gupta<sup>3</sup>, Govind Rankawat<sup>1</sup>

## Abstract

The pandemic of COVID-19 has afflicted every individual and has initiated a cascade of directly or indirectly involved events in precipitating mental health issues. The human species is a wanderer and hunter-gatherer by nature, and physical social distancing and nationwide lockdown have confined an individual to physical isolation. The present review article was conceived to address psychosocial and other issues and their aetiology related to the current pandemic of COVID-19. The elderly age group has most suffered the wrath of SARS-CoV-2, and social isolation as a preventive measure may further induce mental health issues. Animal model studies have demonstrated an inappropriate interacting endogenous neurotransmitter milieu of dopamine, serotonin, glutamate, and opioids, induced by social isolation that could probably lead to observable phenomena of deviant psychosocial behavior. Conflicting and manipulated information related to COVID-19 on social media has also been recognized as a global threat. Psychological stress during the current pandemic in frontline health care workers, migrant workers, children, and adolescents is also a serious concern. Mental health issues in the current situation could also be induced by being quarantined, uncertainty in business, jobs, economy, hampered academic activities, increased screen time on social media, and domestic violence incidences. The gravity of mental health issues associated with the pandemic of COVID-19 should be identified at the earliest. Mental health organization dedicated to current and future pandemics should be established along with Government policies addressing psychological issues to prevent and treat mental health issues need to be developed.

**Keywords:** Psychosocial, Mental Health, COVID-19, Social Isolation, Quarantine, Infodemic

## Background

The pandemic of COVID-19 that began from Wuhan, a city of Hubei province of China, spread like wildfire across every corner of the world with cumulative cases surpassing 23 million and what makes the situation more worrisome is the mortality due to COVID-19 that amounts to 800 906 deaths as of 23 August 2020 worldwide, threatening the very existence of mankind [1]. The current pandemic has afflicted every individual and sector directly or indirectly, and one aspect that has not been addressed appropriately till now is its psychosocial and other consequences that have begun to manifest in some or the other ways. The governments across many countries worldwide imposed lockdown to curb the community spread of COVID-19, considering its menacing fangs of infectivity.

Moreover, practices such as wearing masks, social distancing have also been extensively promoted. The human species is a wanderer and hunter-gatherer by nature, and physical social distancing and the nationwide lockdown has confined an individual to physical isolation and left no choice other than being connected virtually. The world has already witnessed mental health issues pertaining to such kinds of pandemics and their aftermaths [2,3]. The available literature on previous incidences of outbreaks indicates the occurrence of several mental health issues such as irritability, anxiety, depression, insomnia, anger, loneliness, and even suicidal tendencies [4-8]. Evaluating the community's mental health status in such pandemics needs special attention in the Indian context where social gatherings such as marriages, religious festivals, etc. are frequent and spread across the year [9]. Hence, challenges associated with the curbing of community spread of COVID-19 were consequentially increased. Research has shown that around 2/5 of the population is already subjected to anxiety and depression as a consequence of nationwide lockdowns against COVID-19 [10]. The present review article was conceived with

\*Correspondence: [bhupendra.kool9999@gmail.com](mailto:bhupendra.kool9999@gmail.com)

<sup>2</sup>Department of Physiology, Government Medical College, Barmer, Rajasthan, India

Full list of author information is available at the end of the article

the aim of addressing psychosocial and other issues and their aetiology related to the current pandemic of COVID-19. Mental health needs to be stressed during such critical times with the implementation of better government strategies and policies to deal with its psychological consequences.

The pandemic of COVID-19 has initiated a cascade of directly or indirectly involved events in precipitating mental health issues. The list of events is long that includes social isolation, overburdened health care workers, fear and stigma of COVID-19, uncertainty about the economy, fear of losing jobs and loss in business, increased screen time, especially in children and adolescents, misinformation circulating about the pandemic, increased domestic violence and other myriad inconceivable issues as of this date.

### Social isolation in elderly individuals

Social deprivation in an individual's life can have serious mental health consequences. The ongoing pandemic of COVID-19 has compelled the society to remain in social isolation, with a silver lining of being connected virtually through online platforms. Hawkey and Cacioppo (2010) had stressed the importance of the feeling of being socially connected in an individual's life, and lack of this feeling can have deleterious effects on physical and mental health [11]. The elderly age group, in particular, needs special mention, as it has been reported to be the most vulnerable group to have suffered the wrath of SARS-CoV-2 [12,13]. The current situation has drastically impaired the physical, social connectivity, an essential component of innate human needs [14]. The resulting loneliness from social isolation can lead to a greater risk of morbidity and mortality among elderly individuals [15-19]. At this age, individuals are already vulnerable to loneliness, which could be due to any illnesses, death of a spouse, or lack of friends [20]. The social distancing advisory by the government, specially targeted to protect elderly individuals from COVID-19, adds on to already existing loneliness, consequently increasing the risk of mental health issues [21].

Researchers have tried to identify the brain areas related to emotion processing and socialization, and the effects of prolonged social isolation on normal brain functioning [need further elucidation]. Various animal model studies have tried to address this question in entirety. Amygdala has been identified as a seat of social functioning in an individual [22]. Kanai et al. [23] assessed the amygdala's grey matter volume and found a correlation between physical and online social networking and the volume of the amygdala's grey matter. Prolonged social deprivation induces stress that culminates in hippocampal injury due to raised glucocorticoid levels and blood pressure [24-27]. Social isolation has also been recognized as an important risk factor for cognitive dysfunction, such as Alzheimer's disease, where this risk was doubled in lonely individuals [28]. Even an acute social isolation period could cause long-term memory dysfunction, as was evident from mice studies [29].

Various neurotransmitter system dysfunctions have also been demonstrated in experimental studies underlying social isolation in rodents [30-32]. The neurotransmitters dopamine and the serotonin were found to increase in the brain's prefrontal region in experimental mice study, which could underly abnormal behavior due to social isolation [33]. Serotonin has also been implicated in inducing aggression and

depression in a socially isolated animal model [34]. Glutamate levels are also elevated in socially isolated rats, where glutamate-induced excitotoxicity has been demonstrated, subsequently causing dysfunction of receptors and neuronal injury [35, 36]. The resultant brain injury could be attributed to the increased glutamatergic transmission through activation of N-methyl-D-aspartate (NMDA) receptors [37, 38]. These receptors have been found to be involved in the pathophysiology of various mental disorders such as depression, anxiety, and schizophrenia [39-41]. Glutamate receptors have also been reduced under the effect of social isolation in the cortex and hippocampus [42,43]. Furthermore, NMDA receptors' increased binding capacity under social isolation stress in the frontal cortex has also been reported [44]. AMPA ( $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid) is another ionotropic glutamate receptor, whose upregulation and activation in the amygdala could induce anxiety and depression-like behaviors in socially isolated mice [45, 46].

Social isolation also affects the noradrenaline system, where elevated levels have been demonstrated in the rat model's hippocampus, cortex, and cerebellum [47]. It could also lead to aggression induced adrenaline stress response [48]. The opioidergic system is involved in animal studies socialization, which could also be disrupted by social isolation [49-51]. The cannabinoid receptor system is also a vital neuromodulator in the central nervous system, whose depression has been demonstrated to be affected by fear, stress [52, 53]. In a socially isolated mouse model, disruption of the cannabinoid system-induced depression-like picture has been reported [54]. Nitric oxide is another signaling molecule suggested to have a role in learning, memory, synaptic plasticity, stress, seizure susceptibility, pain modulation, and depression in various brain regions [55-58]. Its social isolation role is evident in the studies demonstrating an exhibition of altered modulation of NMDA receptors and nitrenergic system balance in cortico-limbic areas in socially isolated mice [59, 60]. Both systems have also been linked to higher seizure susceptibility in socially deprived rats [35]. Haj-Mirzaian et al. [61] reported that four weeks of juvenile social isolation stress in mice could induce depression and anxiety-like pictures associated with increased hypothalamic responsiveness–pituitary–adrenal (HPA) axis, interleukin-1 $\beta$  upregulation. They increased the production of NO in the prefrontal cortex and hippocampus regions [61]. In conclusion, an inappropriate interacting endogenous neurotransmitter milieu of dopamine, serotonin, glutamate, and opioids could probably lead to observable phenomena of deviant psychosocial behavior that could plausibly have a deleterious effect on neuroeconomics of an individual with dysfunctional decision making distributed across and over time.

### The "Infodemic" alongside Pandemic

The current pandemic of SARS-CoV-2 is accompanied by an "Infodemic" of misinformation and myths are driven due to the consequential response of fear, stigma, and blame-game. The World Health Organization (WHO) has also raised concern about this deluge of conflicting and manipulated information on social media, that poses a global public health threat [62, 63]. The consequences of such an Infodemic could be appreciated by the misinformation spread during the outbreak of Ebola in

the Democratic Republic of Congo in 2019, which stirred violent behavior, lack of trust, and episodes of targeted attacks against health care workers [64]. Reports of similar incidences have surfaced in the news from India and other parts of the world, where health care workers engaged in COVID-19 sample collection and management were assaulted or abused [65-68]. Such kind of behavior, if not tackled timely by releasing correct information and appropriate Government intervention, could impede the efforts being made to manage and curb the spread of COVID-19. Misinformation could also be associated with the stigmatization of people with or suspected of having COVID-19, where such persons may not report the actual status of illness, and subsequently contribute to community spread of the disease [69,70]. Several incidences have been reported from many countries, where due to misinformation, people have suffered serious consequences, such as drinking methanol to cure COVID-19, which led to hospitalization, blindness, or deaths [71-74]. The government must take appropriate measures to track and prevent the spread of misinformation, rumors, stigma, and conspiracy theories on social media platforms, as it can lead to severe implications on people and the community [75].

#### **Psychological stress during the quarantine period**

People quarantined during previous outbreaks have reported a high level of mental stress [76]. The most common complaints include low mood followed by irritability and insomnia [77]. Other symptoms, such as emotional disturbance, [78] depression [76], post-traumatic stress symptoms [79], and anger [80], have also been reported. Health care workers were more susceptible to mental health issues than the general public during the quarantine period, with increased post-traumatic stress severity [79]. Even after three post quarantine years, the hospital staff reported high depressive symptoms [81]. It was also suggested that a quarantine period of more than ten days could lead to higher post-traumatic stress symptoms [76]. The Indian Government has released advisory from time to time on home and institutional quarantine, where more than ten days of quarantine for mild/pre-symptomatic/asymptomatic cases was recommended [82-85]. Although it was necessary to prevent the spread of infection, such preventive measures could create psychological stress in an individual and can have long term consequences affecting the individual and the entire public health system [4]. Hence, there is an urgent need to implement appropriate steps to address related mental health issues.

#### **Business and Economy**

The systemic lockdowns to curb the COVID-19 spread worldwide have significantly hit the economy, that also is a matter of great concern as it will consequently affect the business, entertainment, hotel sectors, and jobs. This pandemic has disrupted significant industry sectors, eventually leading to the closure of many business start-ups. Furthermore, the projected GDP growth is estimated low for the year 2020-12 [86]. This prevailing uncertainty of the economy will further escalate the anxiety and stress in people. There have been fewer incidences of such widespread pandemics, although available literature indicates subsequent low returns on assets [87]. The prevailing uncertainty and fear due to the current pandemic have made us realize that the world was completely unprepared

for such large-scale outbreaks. The loss to the economy will undoubtedly create psychological stress among people, leading to mental health issues.

#### **Psychological stress in migrant workers**

An important issue that needs attention has been of migration of daily wage laborers during the COVID-19 pandemic. This cohort of people was already at risk of psychological stress, as has been reported by Firdaus. They highlighted that mental health issues could pose a significant problem among single, illiterate daily wage laborers, migrant workers lacking appropriate housing, and sanitation [88]. The nation-wide lockdown has resulted in the loss of earnings and jobs, insecurities, and social isolation that is detrimental to the migrant population's mental health. This population was already vulnerable to risk-seeking behavior, as evidenced by the National Crime Records Bureau report, 2018, where 22.4% of suicides among daily wage earners were reported [89]. Another issue that confounded to the prevailing situation was the non-availability of transport to return home amid lockdown, in such a crisis of COVID-19 resulted in significant deaths of migrant workers in road accidents. [90] Such incidences might impact migrant workers' lives and mental health in the long term, possibly leading to high anxiety, depression, and suicidal tendencies. Consequently, social isolation, along with severely compromised available resources, confound further to the burning medical issue of COVID-19.

#### **Children and Adolescent Mental Health**

The COVID-19 seems to have spared the paediatric age group quantitatively, and a relatively milder presentation of the disease has been observed [13, 91]. However, the paediatric age group is more vulnerable to mental health disorders, owing to the confinement itself and an increased screen time due to excessive social media use and online teaching schedule. Additionally, misinformation, exposure to mass media coverage at an earlier age, fear, and stigma of COVID-19 might lead to mental distress in this age group [92-93]. This can be further aggravated by the quarantine of children diagnosed with or suspected of having COVID-19, leading to high levels of anxiety, stress, and even adjustment disorders [94]. At a younger age, such kind of social deprivation, separation from parents, and fear of unknown disease might have a long-lasting psychological impact on children and an even higher risk of developing post-traumatic stress disorder (PTSD) [4]. Previous studies have reported the development of PTSD in 30% of children after being quarantined amid health-related disasters [95].

#### **Education**

The current pandemic has affected various sectors of society directly or indirectly in unexpected ways, including education. The research has shown that students and healthcare professionals are the most vulnerable group developing anxiety, stress, and depression amid COVID-19 lockdown [96]. An unexpected shift in students' learning methods from traditional to virtual classroom teaching has created undue stress on students and teachers. Both have struggled while accessing these online platforms. Teachers might find it challenging to

deal with online teaching platforms and lack of resources efficiently.

In contrast, students might also experience difficulties accessing the online platforms and financial constraints, causing an inability to access the Internet and the unavailability of electronic tools, including the laptops, the phones, the computers, or even the radio and television. Moreover, the lack of adequate and effective interaction during sessions of teaching and knowledge dissemination has resulted in their subsequent pitfalls. However, both government and non-government organizations and educational technology companies are being made efforts to support the school system to make a smooth transition to the virtual world. This altogether has caused high anxiety among students amid the COVID-19 pandemic, together with poor economic conditions and hampered academic activities [97]. The uncertainty and fear of getting infected due to SARS-COV-2 have led to higher university students' anxiety [98].

### Psychological Stress in Health Care Workers

Health care workers (HCW) on the frontline, is the most vulnerable population in the current pandemic. Studies have reported and raised the concern of significant mental health disturbances such as depression, anxiety, and insomnia in health care workers engaged in COVID-19 management [99, 100]. The vulnerability of HCWs for depression, anxiety along with post-traumatic stress disorder was also observed during the previous outbreaks of the middle east respiratory syndrome (MERS) and severe acute respiratory syndrome (SARS) [101-104]. The previous outbreak SARS has shown that healthcare providers involved in the management of SARS-affected patients were also prone to stigmatization associated with the disease [105]. Similar stigma has been associated with the pandemic of COVID-19, where reports of doctors' eviction from their homes and incidences of assault have been documented [106, 65]. The overburdened health care system, misinformation, psychological stress, and uncertainty of the current pandemic may lead to and add to mental health issues among healthcare workers.

Violence against women has also increased globally amid lockdown during this pandemic [107]. Previous studies have also indicated a rise in domestic violence cases during natural disasters [108]. Another concern related to COVID-19 is the increase in screen time during the lockdown, especially in children and adolescents. In present times of social isolation and deprivation with minimal physical contact stimulation and illogical infatuation with social media, it has seemingly made matters worse. There have also been reports of suicide attempts made by people experiencing substance withdrawal [109, 110]. Compliance with forced home quarantine is often being violated in India, unlike in other countries [111].

### Measures for combating psychosocial stress due to COVID-19

There is a need to understand the psychological impact of COVID-19. This will help people across all strata of the society to cope up with the challenges faced. This pandemic could be the much-needed wake-up call to the necessity of long-term changes to India's health system. The psychological stress, anxiety, fear, and helplessness associated with the current

pandemic of COVID-19 needs timely measures to improve society's mental health [112-114]. The disruption of the health care system, including mental health services, [at the time of need when psychological stress in the general population has increased], leaves people helpless [10, 115]. This could, in turn, precipitate incidences of panic, agitation, violence, and increased suicide rates and could be more damaging than the virus itself. The problem should be identified at the earliest, and timely measures targeting psychological issues can prevent crises in such a stressful situation [116].

The previous outbreak of SARS made us realized that psychiatrists could play a critical role in improving the overall health-care service utilization effectively [3]. They may offer a balanced perspective with adequate counselling sessions to provide awareness and mental health education as well as effective management of anxiety and apprehension. The government should intervene early and develop prevention and intervention models targeted at mental health disorders associated with the pandemic. The mental organization targeted at current and future pandemics should also be established [117]. COVID-19 has created mayhem and misery, but there are some positive aspects about lockdowns as well, such as a decrease in pollution level [118], a chance to reconnect with family members [119], and improved family relations and dynamics [120].

### Conclusion

The pandemic of COVID-19 has afflicted every individual and has initiated a cascade of directly or indirectly involved events in precipitating mental health issues. The psychosocial aspect of this pandemic should not be disregarded as it may culminate into various psychiatric disorders. This study presents the various underlying factors that need to be addressed urgently. Governments should intervene early and develop prevention and intervention models targeted at mental health disorders associated with the pandemic. Mental health organization dedicated to current and future pandemics should be established, especially targeting the vulnerable cohort of the society like pediatric, elderly, and frontline workers.

### Abbreviations

SARS-CoV-2: Severe Acute Respiratory Infection-Coronavirus-2; NMDA: N-methyl-D-aspartate; AMPA:  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole Propionic Acid; HPA: Hypothalamic-Pituitary-Adrenal; WHO: World Health Organization; PTSD: Post-Traumatic Stress Disorder; HCW: Health Care Worker; MERS: Middle East Respiratory Syndrome.

### Declarations

### Acknowledgment

None

### Funding

The authors received no financial support for their research, authorship, and/or publication of this article.

### Availability of data and materials

Data will be available by emailing bhupendra.kool9999@gmail.com

### Authors' contributions



BP is the principal investigator of this manuscript (Review), who designed the study and coordinated all aspects of the research, including all steps of the manuscript preparation. He is responsible for the study concept, design, writing, reviewing, editing, and approving the manuscript in its final form. SB, AJS, and AD contributed to the study design, analysis, and reviewed and approved the manuscript. SK, AT, JG and GR contributed to the interpretation of data, drafting the work, writing the manuscript and reviewed and approved the manuscript. All authors read and approved the final manuscript.

### Ethics approval and consent to participate

We conducted the research following the Declaration of Helsinki. However, Review Articles need no ethics committee approval.

### Consent for publication

Not applicable

### Competing interest

The authors declare that they have no competing interests.

### Open Access

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

### Author details

<sup>1</sup>Department of Medicine, S. M. S. Medical College and Attached Hospitals, Jaipur, India. <sup>2</sup>Department of Physiology, Government Medical College, Barmer, Rajasthan, India. <sup>3</sup>Department of Physiology, S. M. S. Medical College and Attached Hospitals, Jaipur, India. <sup>4</sup>Department of Pharmacology, S. M. S. Medical College and Attached Hospitals, Jaipur, India.

### Article Info

Received: 10 September 2020

Accepted: 22 September 2020

Published: 01 October 2020

### References

- World Health Organization (WHO) Coronavirus Disease (COVID-19) Dashboard. Available at: <https://covid19.who.int/> [Accessed on 23 August 2020]
- Sim K, Chua HC. The psychological impact of SARS: a matter of heart and mind. *CMAJ*. 2004; 170:811e2. <https://doi.org/10.1503/cmaj.1032003>.
- Wu P, Fang Y, Guan Z, Fan B, Kong J, Yao Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can J Psychiatr*. 2009; 54:302e11. <https://doi.org/10.1177/070674370905400504>.
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020; 395:912e20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8).
- Robertson E, Hershenfield K, Grace SL, Stewart DE. The psychosocial effects of being quarantined following exposure to SARS: a qualitative study of Toronto health care workers. *Can J Psychiatr*. 2004; 49:403e7. <https://doi.org/10.1177/070674370404900612>.
- Barbisch D, Koenig KL, Shih FY. Is there a case for quarantine? Perspectives from SARS to Ebola. *Disaster Med Public Health Prep*. 2015; 9:547e53. <https://doi.org/10.1017/dmp.2015.38>.
- Jeong H, Yim HW, Song YJ, Ki M, Min JA, Cho J, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health*. 2016;38: e2016048. <https://doi.org/10.4178/epih.e2016048>.
- Liu X, Kakade M, Fuller CJ, Fan B, Fang Y, Kong J, et al. Depression after exposure to stressful events: lessons learned from the severe acute respiratory syndrome epidemic. *Compr Psychiatr*. 2012; 53:15e23. <https://doi.org/10.1016/j.comppsy.2011.02.003>.
- Chadda RK, Deb KS. Indian family systems, collectivistic society and psychotherapy. *Indian J Psychiatry*. 2013;55: S299-309. <https://dx.doi.org/10.4103/2F0019-5545.105555>.
- Grover S, Sahoo S, Mehra A, Avasthi A, Tripathi A, Subramanyan A, et al. Psychological impact of COVID-19 lockdown: An online survey from India. *Indian J Psychiatry*. 2020; 62:354-62. [https://doi.org/10.4103/psychiatry.IndianJPsychiatry\\_427\\_20](https://doi.org/10.4103/psychiatry.IndianJPsychiatry_427_20).
- Hawkey LC, Cacioppo JT. Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Ann Behav Med*. 2010; 40: 218–27. <https://dx.doi.org/10.1007/2Fs12160-010-9210-8>.
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-13. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7).
- Bhandari S, Sharma R, Singh Shaktawat A, Banerjee S, Patel B, Tak A, et al. COVID-19 related mortality profile at a tertiary care centre: a descriptive study. *Scr Med*. 2020;51(2):69-73. <https://doi.org/10.5937/scriptamed51-27126>.
- Baumeister RF, Leary MR. The need to belong: desire for interpersonal attachments as a fundamental human motivation. *Psychol Bull*. 1995; 117: 497–529. <https://doi.org/10.1037/0033-2909.117.3.497>.
- Caspi A, Harrington H, Moffitt TE, Milne BJ, Poulton R. Socially isolated children 20 years later: risk of cardiovascular disease. *Arch Pediatr Adolesc Med*. 2006; 160(8):805-11. <https://doi.org/10.1001/archpedi.160.8.805>.
- Eaker ED, Pinsky J, Castelli WP. Myocardial infarction and coronary death among women: psychosocial predictors from a 20-year follow-up of women in the Framingham Study. *Am J Epidemiol*. 1992; 135(8):854-64. <https://doi.org/10.1093/oxfordjournals.aje.a116381>.
- Luo Y, Hawkey LC, Waite LJ, Cacioppo JT. Loneliness, health, and mortality in old age: a national longitudinal study. *Soc Sci Med*. 2012 Mar; 74(6):907-14. <https://dx.doi.org/10.1016%2Fj.socscimed.2011.11.028>.
- Olsen RB, Olsen J, Gunner-Svensson F, Waldstrøm B. Social networks and longevity. A 14-year follow-up study among elderly in Denmark. *Soc Sci Med*. 1991; 33(10):1189-95. [https://doi.org/10.1016/0277-9536\(91\)90235-5](https://doi.org/10.1016/0277-9536(91)90235-5).
- Patterson AC, Veenstra G. Loneliness and risk of mortality: a longitudinal investigation in Alameda County, California. *Soc Sci Med*. 2010; 71(1):181-6. <https://doi.org/10.1016/j.socscimed.2010.03.024>.
- Savikko N, Routassalo P, Tilvis RS, Strandberg TE, Pitkalla KH. Predictors and subjective causes of loneliness in an aged population. *Arch Gerontol Geriatrics*. 2005; 41:3:223-33. <https://doi.org/10.1016/j.archger.2005.03.002>.
- Health Advisory for Elderly Population of India during COVID19. Available at: <https://www.mohfw.gov.in/pdf/AdvisoryforElderlyPopulation.pdf> [Accessed on 13 August 2020].
- Dicks D, Myers R, Kling A. Uncus and amygdala lesions: effects on social behavior in the free-ranging rhesus monkey. *Science*. 1969; 165:69–71. <https://doi.org/10.1126/science.165.3888.69>.
- Kanai R, Bahrami B, Duchaine B, Janik A, Banissy MJ, Rees G. Brain structure links loneliness to social perception. *Curr Biol*.

- 2012; 22(20):1975-9. <https://dx.doi.org/10.1016%2Fj.cub.2012.08.045>.
24. Bender AR, Daugherty A, Raz N. Vascular risk moderates associations between hippocampal subfield volumes and memory. *J Cogn Neurosci*. 2013; 25:1851-62. [https://doi.org/10.1162/jocn\\_a\\_00435](https://doi.org/10.1162/jocn_a_00435).
  25. Raz N. Diabetes: brain, mind, insulin—what is normal and do we need to know? *Nat Rev Endocrinol*. 2011; 7:636-7. <https://doi.org/10.1038/nrendo.2011.149>.
  26. Colcombe SJ, Erickson KI, Naftali R, Andrew GW, Cohen NJ, McAuley E, et al. Aerobic fitness reduces brain tissue loss in aging humans. *J Gerontol A Biol Sci Med Sci*. 2003; 58:176-80. <https://doi.org/10.1093/gerona/58.2.m176>.
  27. Maass A, Düzel S, Goerke M, Becke A, Sobieray U, Neumann K, et al. Vascular hippocampal plasticity after aerobic exercise in older adults. *Mol Psychiatry*. 2015; 20, 585-93. <https://doi.org/10.1038/mp.2014.114>.
  28. Wilson RS, Krueger KR, Arnold SE, Schneider JA, Kelly JF, Barnes LL, et al. Loneliness and Risk of Alzheimer Disease. *Arch Gen Psychiatry*. 2007;64(2):234-240. <https://doi.org/10.1001/archpsyc.64.2.234>.
  29. Kogan JH, Frankland PW, Silva AJ. Long-term memory underlying hippocampus-dependent social recognition in mice. *Hippocampus*. 2000;10(1):47-56. [https://doi.org/10.1002/\(sici\)1098-1063\(2000\)10:1%3C47::aid-hipo5%3E3.0.co;2-6](https://doi.org/10.1002/(sici)1098-1063(2000)10:1%3C47::aid-hipo5%3E3.0.co;2-6).
  30. Yorgason JT, España RA, Konstantopoulos JK, Weiner JL, Jones SR. Enduring increases in anxiety-like behavior and rapid nucleus accumbens dopamine signaling in socially isolated rats. *Eur J Neurosci*. 2013;37(6):1022-31. <https://doi.org/10.1111/ejn.12113>.
  31. Bledsoe AC, Oliver KM, Scholl JL, Forster GL. Anxiety states induced by post-weaning social isolation are mediated by CRF receptors in the dorsal raphe nucleus. *Brain Res Bull*. 2011;85(3-4):117-22. <https://dx.doi.org/10.1016%2Fj.brainresbull.2011.03.003>.
  32. Lukkes JL, Engelman GH, Zelin NS, Hale MW, Lowry CA. Post-weaning social isolation of female rats, anxiety-related behavior, and serotonergic systems. *Brain Res*. 2012; 1443:1-17. <https://dx.doi.org/10.1016%2Fj.brainres.2012.01.005>.
  33. Ago Y, Araki R, Tanaka T, Sasaga A, Nishiyama S, Takuma K, et al. Role of social encounter-induced activation of prefrontal serotonergic systems in the abnormal behaviors of isolation-reared mice. *Neuropsychopharmacology*. 2013; 38(8):1535-47. <https://doi.org/10.1038/npp.2013.52>.
  34. Veenema AH. Early life stress, the development of aggression and neuroendocrine and neurobiological correlates: what can we learn from animal models? *Front Neuroendocrinol*. 2009;30(4):497-518. <https://doi.org/10.1016/j.yfrne.2009.03.003>.
  35. Zhao X, Sun L, Jia H, Meng Q, Wu S, Li N, et al. Isolation rearing induces social and emotional function abnormalities and alters glutamate and neurodevelopment-related gene expression in rats. *Prog Neuropsychopharmacol Biol Psychiatry*. 2009;33(7):1173-1177. <https://doi.org/10.1016/j.pnpbp.2009.06.016>.
  36. Sciolino NR, Bortolato M, Eisenstein SA, Fu J, Oveisi F, Hohmann AG, et al. Social isolation and chronic handling alter endocannabinoid signaling and behavioral reactivity to context in adult rats. *Neuroscience*. 2010;168(2):371-86. <https://dx.doi.org/10.1016%2Fj.neuroscience.2010.04.007>.
  37. Ghasemi M, Phillips C, Trillo L, De Miguel Z, Das D, Salehi A. The role of NMDA receptors in the pathophysiology and treatment of mood disorders. *Neurosci Biobehav Rev*. 2014; 47:336-358. <https://doi.org/10.1016/j.neubiorev.2014.08.017>.
  38. Olivenza R, Moro MA, Lizasoain I, Lorenzo P, Fernández AP, Rodrigo J, et al. Chronic stress induces the expression of inducible nitric oxide synthase in rat brain cortex. *J Neurochem*. 2000;74(2):785-791. <https://doi.org/10.1046/j.1471-4159.2000.740785.x>.
  39. Maeng S, Zarate CA Jr, Du J, Schloesser RJ, McCammon J, Chen G, et al. Cellular mechanisms underlying the antidepressant effects of ketamine: role of alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptors. *Biol Psychiatry*. 2008;63(4):349-352. <https://doi.org/10.1016/j.biopsych.2007.05.028>.
  40. Kalia LV, Kalia SK, Salter MW. NMDA receptors in clinical neurology: excitatory times ahead. *Lancet Neurol*. 2008;7(8):742-755. [https://dx.doi.org/10.1016%2FS1474-4422\(08\)70165-0](https://dx.doi.org/10.1016%2FS1474-4422(08)70165-0).
  41. Waxman EA, Lynch DR. N-methyl-D-aspartate Receptor Subtypes: Multiple Roles in Excitotoxicity and Neurological Disease. *The Neuroscientist*. 2005; 11(1), 37-49. <https://doi.org/10.1177/1073858404269012>.
  42. Hermes G, Li N, Duman C, Duman R. Post-weaning chronic social isolation produces profound behavioral dysregulation with decreases in prefrontal cortex synaptic-associated protein expression in female rats. *Physiol Behav*. 2011;104(2):354-9. <https://dx.doi.org/10.1016%2Fj.physbeh.2010.12.019>.
  43. Sestito RS, Trindade LB, de Souza RG, Kerbauy LN, Iyomasa MM, Rosa ML. Effect of isolation rearing on the expression of AMPA glutamate receptors in the hippocampal formation. *J Psychopharmacol*. 2011;25(12):1720-1729. <https://doi.org/10.1177/0269881110385595>.
  44. Toua C, Brand L, Möller M, Emsley RA, Harvey BH. The effects of sub-chronic clozapine and haloperidol administration on isolation rearing induced changes in frontal cortical N-methyl-D-aspartate and D1 receptor binding in rats. *Neuroscience*. 2010;165(2):492-499. <https://doi.org/10.1016/j.neuroscience.2009.10.039>.
  45. Alò R, Avolio E, Mele M, Storino F, Canonaco A, Carelli A et al. Excitatory/inhibitory equilibrium of the central amygdala nucleus gates anti-depressive and anxiolytic states in the hamster. *Pharmacol Biochem Behav*. 2014; 118:79-86. <https://doi.org/10.1016/j.pbb.2014.01.007>.
  46. St JP, Petkov VV. Changes in 5-HT1 receptors in different brain structures of rats with isolation syndrome. *General pharmacology*. 1990;21(2):223-5. [https://doi.org/10.1016/0306-3623\(90\)90905-2](https://doi.org/10.1016/0306-3623(90)90905-2).
  47. Miachon S, Rochet T, Mathian B, Barbagli B, Claustrat B. Long-term isolation of Wistar rats alters brain monoamine turnover, blood corticosterone, and ACTH. *Brain Res Bull*. 1993;32(6):611-614. [https://doi.org/10.1016/0361-9230\(93\)90162-5](https://doi.org/10.1016/0361-9230(93)90162-5).
  48. Van den Berg CL, Van Ree JM, Spruijt BM, Kitchen I. Effects of juvenile isolation and morphine treatment on social interactions and opioid receptors in adult rats: behavioural and autoradiographic studies. *Eur J Neurosci*. 1999;11(9):3023-3032. <https://doi.org/10.1046/j.1460-9568.1999.00717.x>.
  49. Vanderschuren LJ, Stein EA, Wiegant VM, Van Ree JM. Social play alters regional brain opioid receptor binding in juvenile rats. *Brain Res*. 1995;680(1-2):148-156. [https://doi.org/10.1016/0006-8993\(95\)00256-p](https://doi.org/10.1016/0006-8993(95)00256-p).
  50. Moles A, Kieffer BL, D'Amato FR. Deficit in attachment behavior in mice lacking the mu-opioid receptor gene. *Science*. 2004;304(5679):1983-1986. <https://doi.org/10.1126/science.1095943>.
  51. Panksepp J, Herman BH, Vilberg T, Bishop P, DeEsquinazi FG. Endogenous opioids and social behavior. *Neurosci Biobehav Rev*. 1980;4(4):473-487. [https://doi.org/10.1016/0149-7634\(80\)90036-6](https://doi.org/10.1016/0149-7634(80)90036-6).
  52. Gong JP, Onaivi ES, Ishiguro H, Liu Q, Tagliaferro PA, Brusco A, et al. Cannabinoid CB2 receptors: immunohistochemical localization in rat brain. *Brain Res*. 2006;1071(1):10-23. <https://doi.org/10.1016/j.brainres.2005.11.035>.
  53. Breivogel CS, Sim-Selley LJ. Basic neuroanatomy and neuropharmacology of cannabinoids. *Int Rev Psychiatry* 2009; 21:2:113-121. <https://doi.org/10.1080/09540260902782760>.
  54. Haj-Mirzaian A, Amini-Khoei H, Haj-Mirzaian A, Amiri S, Ghesmati M, Zahir M, et al. Activation of cannabinoid receptors elicits antidepressant-like effects in a mouse model of social isolation stress. *Brain Res Bull*. 2017; 130:200-210. <https://doi.org/10.1016/j.brainresbull.2017.01.018>.
  55. Banach M, Piskorska B, Czuczwar SJ, Borowicz KK. Nitric Oxide, Epileptic Seizures, and Action of Antiepileptic Drugs. *CNS & Neurological Disorders - Drug Targets* 2011;10: 808. <https://doi.org/10.2174/187152711798072347>.

56. Förstermann U, Sessa WC. Nitric oxide synthases: regulation and function. *Eur Heart J*. 2012;33(7):829-37, 837a-837d. <https://dx.doi.org/10.1093%2Feurheartj%2Fehf304>.
57. Hu Y, Wu D, Luo C, Zhu L, Zhang J, Wu H, et al. Hippocampal nitric oxide contributes to sex difference in affective behaviors. *PNAS*. 2012, 109 (35) 14224-14229. <https://doi.org/10.1073/pnas.1207461109>.
58. Khan MI, Ostadhadi S, Zolfaghari S, Mehr SE, Hassanzadeh G, Dehpour, A et al. The involvement of NMDA receptor/NO/cGMP pathway in the antidepressant like effects of baclofen in mouse force swimming test. *Neuroscience Letters*. 2016; 612:52-61. <https://doi.org/10.1016/j.neulet.2015.12.006>.
59. Matsumoto K, Puia G, Dong E, Pinna G. GABAA receptor neurotransmission dysfunction in a mouse model of social isolation-induced stress: Possible insights into a non-serotonergic mechanism of action of SSRIs in mood and anxiety disorders. *Stress*. 2007; 10(1):3-12. <https://doi.org/10.1080/10253890701200997>.
60. Zlatković J, Filipović D. Chronic social isolation induces NF-κB activation and upregulation of iNOS protein expression in rat prefrontal cortex. *Neurochem Int*. 2013;63(3):172-179. <https://doi.org/10.1016/j.neuint.2013.06.002>.
61. Haj-Mirzaian A, Amiri S, Kordjazy N, Momeny M, Razmi A, Balaei MR, et al. Lithium attenuated the depressant and anxiogenic effect of juvenile social stress through mitigating the negative impact of interleukin-1β and nitric oxide on hypothalamic-pituitary-adrenal axis function. *Neuroscience*. 2016; 315:271-285. <https://doi.org/10.1016/j.neuroscience.2015.12.024>.
62. Larson HJ. The biggest pandemic risk? Viral misinformation. *Nature* 2018; 562:309. <https://doi.org/10.1038/d41586-018-07034-4>.
63. Zarocostas J. How to fight an infodemic. *Lancet* 2020; 395:676. [https://doi.org/10.1016/S0140-6736\(20\)30461-X](https://doi.org/10.1016/S0140-6736(20)30461-X).
64. World Health Organization, 2019. Ebola Virus Disease – Democratic Republic of the Congo. Geneva, Switzerland: WHO. Available at: <https://www.who.int/csr/don/28-november-2019-ebola-drc/en/> [Accessed on August 8, 2020]
65. Times of India. Covid-19: doctors gone to collect samples attacked in Indore. Available at: <https://timesofindia.indiatimes.com/videos/news/covid-19-doctors-goneto-collect-samples-attacked-in-indore/videoshow/74942153.cms; 2020> [Accessed on August 8, 2020].
66. Withnall A. Coronavirus: why India has had to pass new law against attacks on healthcare workers. *The Independent*. April 23, 2020.
67. Semple K. “Afraid to be a nurse”: health workers under attack. *The New York Times*. 2020 Apr 27.
68. The Economist. Health workers become unexpected targets during COVID-19. *The Economist*. May 11, 2020.
69. Turan B, Budhwani H, Fazeli PL, Browning WR, Raper JL, Mugavero MJ, et al. How does stigma affect people living with HIV? The mediating roles of internalized and anticipated HIV stigma in the effects of perceived community stigma on health and psychosocial outcomes. *AIDS Behav*. 2017; 21: 283–291. <https://doi.org/10.1007/s10461-016-1451-5>.
70. James PB, Wardle J, Steel A, Adams J. An assessment of Ebola-related stigma and its association with informal healthcare utilisation among Ebola survivors in Sierra Leone: a cross sectional study. *BMC Public Health*. 2020; 20: 182. <https://doi.org/10.1186/s12889-020-8279-7>.
71. Aljazeera, 2020. Iran: Over 700 Dead after Drinking Alcohol to Cure Coronavirus. *Aljazeera*. Available at: <https://www.aljazeera.com/news/2020/04/iran-700-dead-drinking-alcohol-cure-coronavirus200427163529629.html>. (Accessed June 4, 2020)
72. Delirrad M, Mohammadi AB, 2020. New methanol poisoning outbreaks in Iran following COVID-19 pandemic. *Alcohol Alcohol*. 55: 347–348. <https://doi.org/10.1093/alcal/agaa036>.
73. Hassanian-Moghaddam H, Zamani N, Kolahi A-A, McDonald R, Hovda KE. Double trouble: methanol outbreak in the wake of the COVID-19 pandemic in Iran—a cross-sectional assessment. *Crit Care*. 2020; 24: 402. <https://doi.org/10.1186/s13054-020-03140-w>.
74. Soltaninejad K. Methanol Mass Poisoning Outbreak: A Consequence of COVID-19 Pandemic and Misleading Messages on Social Media. *Int J Occup Environ Med*. 2020;11(3):148-150. <https://dx.doi.org/10.34172%2Fijoem.2020.1983>.
75. Islam MS, Sarkar T, Khan SH, Kamal AM, Hasan SMM, Kabir A, et al. COVID-19–Related Infodemic and Its Impact on Public Health: A Global Social Media Analysis. *Am J Trop Med Hyg*. 2020; 00(0):1–9. <https://doi.org/10.4269/ajtmh.20-0812>.
76. Hawryluck L, Gold W, Robinson S, Pogorski S, Galea S, Styr R. SARS control and psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis*. 2004;10(7):1206–1212. <https://dx.doi.org/10.3201%2Fid1007.030703>.
77. Lee S, Chan LYY, Chau AAM, Kwok KPS, Kleinman A. The experience of SARS-related stigma at Amoy Gardens. *Soc Sci Med*. 2005; 61(9): 2038-2046. <https://doi.org/10.1016/j.socscimed.2005.04.010>.
78. Yoon MK, Kim SY, Ko HS, Lee MS. System effectiveness of detection, brief intervention and refer to treatment for the people with post-traumatic emotional distress by MERS: a case report of community-based proactive intervention in South Korea. *Int J Ment Health Syst*. 2016; 10: 51. <https://doi.org/10.1186/s13033-016-0083-5>.
79. Reynolds DL, Garay JR, Deamond SL, Moran MK, Gold W, Styr R. Understanding, compliance and psychological impact of the SARS quarantine experience. *Epidemiol Infect*. 2008; 136: 997-1007. <https://dx.doi.org/10.1017%2FS0950268807009156>.
80. Marjanovic Z, Greenglass ER, Coffey S. The relevance of psychosocial variables and working conditions in predicting nurses' coping strategies during the SARS crisis: an online questionnaire survey. *Int J Nurs Stud*. 2007; 44(6): 991-998. <https://doi.org/10.1016/j.ijnurstu.2006.02.012>.
81. Bai Y, Lin C-C, Lin C-Y, Chen J-Y, Chue C-M, Chou P. Survey of stress reactions among health care workers involved with the SARS outbreak. *Psychiatr Serv*. 2004; 55: 1055-1057. <https://doi.org/10.1176/appi.ps.55.9.1055>.
82. Ministry of Health and Family Welfare. Available at: <https://www.mohfw.gov.in/pdf/Guidelinesforhomequarantine.pdf> [Accessed on 25 August 2020].
83. Ministry of Health and Family Welfare. Available at: <https://www.mohfw.gov.in/pdf/RevisedguidelinesforHomeIsolationofverymildpresymptomaticCOVID19cases10May2020.pdf> [Accessed on 25 August 2020].
84. Ministry of Health and Family Welfare. Available at: <https://www.mohfw.gov.in/pdf/AdvisoryformanagingHealthcareworkersworkinginCOVIDandNonCOVIDareasofthehospital.pdf> (Accessed on 25 August 2020).
85. Ministry of Health and Family Welfare. Available at: <https://www.mohfw.gov.in/pdf/RevisedguidelinesforInternationalArrivals02082020.pdf> [Accessed on 25 August 2020].
86. Cost of the lockdown? Over 10% of GDP loss for 18 states. Available at: <https://timesofindia.indiatimes.com/india/cost-of-the-lockdown-over-10-of-gdp-loss-for-18-states/articleshow/76028826.cms> [Accessed on 21 August 2020].
87. Jorda O, Singh SR, Taylor AM. Longer-Run Economic Consequences of Pandemics. *Federal Reserve Bank of San Francisco Working Paper*. 2020-09. <https://doi.org/10.24148/wp2020-09>.
88. Firdaus G. Mental well-being of migrants in urban center of India: Analyzing the role of social environment. *Indian J Psychiatry*. 2017; 59:164-9. [https://doi.org/10.4103/psychiatry.indianjpsychiatry\\_272\\_15](https://doi.org/10.4103/psychiatry.indianjpsychiatry_272_15).
89. National Crime Record Bureau. *Annual Crime in India Report*. New Delhi, India: Ministry of Home Affairs; 2018.
90. 198 migrant workers killed in road accidents during lockdown: Report. Available at: <https://www.hindustantimes.com/india-news/198-migrant-workers-killed-in-road-accidents-during-lockdown-report/story-hTWzAWMYn0kyyKw1dyKqL.html> [Accessed on 25 August 2020].
91. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease



- 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis.* 2020; 20:689-96. [https://doi.org/10.1016/S1473-3099\(20\)30198-5](https://doi.org/10.1016/S1473-3099(20)30198-5).
92. Dalton L, Rapa E, Stein A. Protecting the psychological health of through effective communication about COVID-19. *Lancet Child Adolesc Health.* 2020;4(5):346-347. [https://doi.org/10.1016/S2352-4642\(20\)30097-3](https://doi.org/10.1016/S2352-4642(20)30097-3).
  93. Centre for Disease Control. Helping Children Cope with Emergencies. Available at: <https://www.cdc.gov/childrenindisasters/helping-children-cope.html> [Accessed on 25 August 2020].
  94. Liu JJ, Bao Y, Huang X, Shi J, Lu L. Mental health considerations for children quarantined because of COVID-19. *Lancet Child & Adolesc Health.* 2020; 4(5):347-349. [https://doi.org/10.1016/S2352-4642\(20\)30096-1](https://doi.org/10.1016/S2352-4642(20)30096-1).
  95. Sprang G, Silman M. Posttraumatic Stress Disorder in Parents and Youth After Health-Related Disasters. *Disaster Med Public Health Prep.* 2013;7(1):105-110. <https://doi.org/10.1017/dmp.2013.22>.
  96. Rehman U, Shahnawaz MG, Khan NH, Kharshiing KD, Khursheed M, Gupta K, et al. Depression, Anxiety and Stress Among Indians in Times of Covid-19 Lockdown. *Community Ment Health J.* 2020:1-7. <https://doi.org/10.1007/s10597-020-00664-x>.
  97. Cao W, Fang Z, Hou, Han M, Xu X, Dong J, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Research.* 2020; 287:112934. <https://doi.org/10.1016/j.psychres.2020.112934>.
  98. Wang C, Zhao H. The Impact of COVID-19 on Anxiety in Chinese University Students. *Front Psychol.* 2020; 11:1168. <https://dx.doi.org/10.3389/fpsyg.2020.01168>.
  99. Kang L, Li Y, Hu S, Chen M, Yang C, Yang BX, et al. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry* 2020;7(3): e14. [https://doi.org/10.1016/s2215-0366\(20\)30047-x](https://doi.org/10.1016/s2215-0366(20)30047-x).
  100. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open* 2020;3(3): e203976. <https://doi.org/10.1001/jamanetworkopen.2020.3976>.
  101. Lancee WJ, Maunder RG, Goldbloom DS, Coauthors for the Impact of SARS Study. Prevalence of psychiatric disorders among Toronto hospital workers one to two years after the SARS outbreak. *Psychiatr Serv.* 2008;59(1):91-95. <https://dx.doi.org/10.1176%2Fps.2008.59.1.91>.
  102. Tam CWC, Pang EPF, Lam LCW, Chiu HFK. Severe acute respiratory syndrome (SARS) in Hongkong in 2003: Stress and psychological impact among frontline healthcare workers. *Psychol Med.* 2004;34 (7):1197-1204. <https://doi.org/10.1017/s0033291704002247>.
  103. Lee SM, Kang WS, Cho A-R, Kim T, Park JK. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. *Compr Psychiatry.* 2018; 87:123-127. <https://dx.doi.org/10.1016%2Fj.comppsy.2018.10.003>.
  104. Koh D, Meng KL, Chia SE, Ko SM, Qian F, Ng V, et al. Risk perception and impact of severe acute respiratory syndrome (SARS) on work and personal lives of healthcare workers in Singapore: What can we learn? *Med Care.* 2005;43(7):676-682. <https://doi.org/10.1097/01.mlr.0000167181.36730.cc>.
  105. Verma S, Mythily S, Chan YH, Deslypere JP, Teo EK, Chong SA. Post-SARS psychological morbidity and stigma among general practitioners and traditional Chinese medicine practitioners in Singapore. *Ann Acad Med Singap.* 2004; 33(6):743e8.
  106. Yeung J, Gupta S. Doctors evicted from their homes in India as fear spreads amid coronavirus lockdown. *CNN World.* 2020. Available at: <https://edition.cnn.com/2020/03/25/asia/india-coronavirus-doctors-discrimination-intl-hnk/index.html>. [Accessed on 24 August 2020]
  107. Violence Against Women and Girls: the Shadow Pandemic. UN Women. 2020. May 3, 2020. Available at: <https://www.unwomen.org/en/news/stories/2020/4/statement-ed-phumzile-violence-against-women-during-pandemic>. [Accessed on 24 August 2020].
  108. Gearhart S, Patron MP, Hammond TA, Goldberg DW, Klein A, Horney JA. The impact of natural disasters on domestic violence: an analysis of reports of simple assault in Florida (1999–2007). *Violence Gen.* 2018;5(2):87–92. <https://doi.org/10.1089/vio.2017.0077>.
  109. Sahoo S, Rani S, Parveen S, Pal Singh A, Mehra A, Chakrabarti S, et al. Self-harm and COVID-19 pandemic: An emerging concern – A report of 2 cases from India. *Asian J Psychiatr* 2020; 51:102104. <https://dx.doi.org/10.1016%2Fj.ajp.2020.102104>.
  110. Ghosh A, Khitiz MT, Pandiyan S, Roub F, Grover S. Multiple suicide attempts in an individual with opioid dependence: Unintended harm of lockdown during the COVID-19 outbreak? *Indian J Psychiatry* 2020; [In Press].
  111. The Economic Times. 11 Coronavirus suspects flee from a hospital in Maharashtra. March 16 2020. Available at: <https://economictimes.indiatimes.com/news/politics-and-nation/11-coronavirus-suspects-flee-from-a-hospital-in-maharashtra/videoshw/74644936.cms?from=mdr>. [Accessed on 23 August 2020].
  112. Xiang Y, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *The Lancet Psychiatry* 2020;(3):228–229. [https://doi.org/10.1016/S2215-0366\(20\)30046-8](https://doi.org/10.1016/S2215-0366(20)30046-8).
  113. Van Bortel T, Basnayake A, Wurie F, Jambai M, Koroma A, Muana A, et al. Psychosocial effects of an Ebola outbreak at individual, community and international levels. *Bull World Health Organ.* 2016;94(3):210–214. <https://dx.doi.org/10.2471%2FBLT.15.158543>.
  114. Kumar A, Nayar KR. COVID 19 and its mental health consequences. *Journal of Mental Health.* 2020; ahead of print:1-2. <https://doi.org/10.1080/09638237.2020.1757052>.
  115. Gupta R, Grover S, Basu A, Krishnan V, Tripathi A, Subramanyam A, et al. Changes in sleep pattern and sleep quality during COVID-19 lockdown. *Indian J Psychiatry.* 2020; 62(4):370-8. [https://doi.org/10.4103/psychiatry.indianjpsychiatry\\_523\\_20](https://doi.org/10.4103/psychiatry.indianjpsychiatry_523_20).
  116. Duan L, Zhu G. Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry.* 2020;7(4): P300-302. [https://doi.org/10.1016/S2215-0366\(20\)30073-0](https://doi.org/10.1016/S2215-0366(20)30073-0).
  117. Dubey S, Biswas P, Ghosh R, Chatterjee S, Dubey MJ, Chatterjee S et al. Psychosocial impact of COVID-19. *Diabetes Metab Syndr.* 2020; 14(5): 779–788. <https://dx.doi.org/10.1016%2Fj.dsx.2020.05.035>.
  118. Wright R. The world's largest coronavirus lockdown is having a dramatic impact on pollution in India. *CNN World;* 2020. Available at: <https://edition.cnn.com/2020/03/31/asia/coronavirus-lockdown-impact-pollution-india-intl-hnk/index.html>. [Accessed on 23 August 2020]
  119. Foster O. 'Lockdown made me Realise What's Important': Meet the Families Reconnecting Remotely. *The Guardian;* 2020. Available at: <https://www.theguardian.com/keep-connected/2020/apr/23/lockdown-made-me-realise-whats-important-meet-the-families-reconnecting-remotely>. (Accessed on 23 August 2020)
  120. Bilefsky D, Yeginsu C. Of 'Covidivorces' and 'Coronababies': Life During a Lockdown. *N. Y. Times;* 2020. Available at: <https://www.nytimes.com/2020/03/27/world/coronavirus-lockdown-relationships.html> [Accessed on 23 August 2020]